



INVITATION

{{accLname}} {{customText[醫師|主任|教授|副院長|院長|護理師|個管師]}} 您好
邀請您參加3月10日晚間7:30舉辦之線上研討會：

How I treat HR+, HER2- breast cancer patients with CDK4&6i

本次邀請劉峻宇醫師及戴明榮醫師與我們分享臨床治療經驗，並由張源清醫師擔任座長帶領與會者進行討論，期待您的參與。

Agenda

- Welcome & Opening - 5 min
- Challenging case: P't with visceral disease - 20 min
- Challenging case: P't with endocrine-resistance - 20 min
- Interactive discussion & exchange - 30 min
 - Treatment algorithm and real world practice
 - Controversial issues
- Wrap up & Closing - 5 min




3/10 (Wed)




19:30 - 20:50

點此完成報名



Moderator

張源清 醫師
台北馬偕醫院



Speaker

劉峻宇 醫師
台北榮民總醫院



Speaker

戴明榮 醫師
三軍總醫院

Sincerely,

{{userName}}

Eli Lilly and Company (Taiwan)

11/F, 365, Fu Hsing N. Road, Taipei, Taiwan 105 R.O.C.

{{userEmailAddress}} www.lilly.com



P-MKT-C4-011-21-Feb-22 | © 2021 Eli Lilly and Company. All Rights Reserved
We respect your right to privacy - [Privacy Statement](#) | [Unsubscribe](#)

CONFIDENTIALITY AND NO FURTHER DISTRIBUTION NOTICE:

You are receiving this email because you are a Healthcare Professional (HCP) registered to receive information from Eli Lilly and Company ("Lilly"). If you are not an HCP or not able to prescribe medicines in Taiwan, please delete and destroy any copies of this email as got to you in error.

This communication is intended for a specific group of HCP only in a restricted group of countries. The content, web links, and/or webinar contained in this email may be promotional in nature and/or may include comprehensive information on Lilly products and/or devices, with an approved indication in Taiwan, and should not be further distributed. Note: Certain web links contained in this email may link to sites operated by other third parties; they are independent from Lilly and Lilly has no control over, or responsibility for, their information or accuracy.

This email message (including all attachments and the content and web links contained therein) are also for the sole use of the intended recipient and may also contain confidential information. Any unauthorized review, use, disclosure, copying, or distribution is strictly prohibited.

If you are not the intended recipient, please contact the sender by reply email and destroy copies of the original message
This email was sent by: Eli Lilly and Company.

11/F, 365, Fu Hsing N. Road Taipei, 105, Taiwan

捷癒寧®膜衣錠 50毫克、100毫克、150毫克、200毫克

VERZENIO® (abemaciclib) film-coated tablet 50mg, 150mg, 200mg

適應症:

- 併用芳香環酶抑制劑(aromatase inhibitor)，可做為治療停經後荷爾蒙受體(HR)陽性、第二型人類表皮生長因子受體(HER2)陰性之晚期或轉移性乳癌婦女的第一線內分泌療法(endocrine-based therapy)。
- 併用fulvestrant，可治療荷爾蒙受體(HR)陽性、第二型人類表皮生長因子受體(HER2)陰性，且接受內分泌療法後疾病惡化之晚期或轉移性乳癌婦女。
- 單獨用於治療荷爾蒙受體(HR)陽性、第二型人類表皮生長因子受體(HER2)陰性，曾經接受過內分泌治療及於轉移後接受化學治療後又發生疾病惡化之晚期或轉移性乳癌的成人病人。

用法用量: 與fulvestrant或芳香環酶抑制劑併用時，VERZENIO的建議劑量為150 mg，每日口服兩次。單獨使用時，VERZENIO的建議劑量為200 mg，每日口服兩次。持續治療直到疾病惡化或出現無法耐受的毒性。VERZENIO可隨餐或空腹使用。

禁忌: 對於VERZENIO內之有效成份或賦形劑過敏者。

警語和注意事項:

腹瀉 病人一旦發現有稀便的情況，應開始止瀉劑治療(如loperamide)、增加口服液體攝取。嗜中性球低下症 VERZENIO治療開始前、治療開始後前2個月每2週一次、之後的2個月每月一次，及當臨床需要時都應監測全血球計數。間質性肺病/肺發炎 監測病人是否出現間質性肺病/肺發炎徵象的肺部症狀。症狀可能包含缺氧、咳嗽、呼吸困難或影像學檢查有間質浸潤。應適當調查以排除感染、腫瘤及其他造成這類症狀的原因。肝毒性 VERZENIO治療開始前及治療開始後的前2個月每2週一次、之後的2個月每月一次，及當臨床需要時都應監測肝功能檢測(LFT)。靜脈血栓栓塞 監測病人是否出現靜脈血栓和肺栓塞的徵候與症狀。胚胎-胎兒毒性 告知懷孕婦女此藥物可能對胎兒造成的潛在危害。告知具有生育能力的女性，在接受VERZENIO治療期間至接受最後一劑VERZENIO治療後至少3週內應採取有效的避孕措施。

不良反應: 最常見之不良反應包括腹瀉、感染、嗜中性球低下症、貧血、疲倦、噁心、嘔吐和食慾不振。

使用前請詳閱VERZENIO®藥品仿單警語及注意事項。

VERZENIO 50毫克 衛部藥輸字第027643號
VERZENIO 100毫克 衛部藥輸字第027642號
VERZENIO 150毫克 衛部藥輸字第027641號
VERZENIO 200毫克 衛部藥輸字第027640號

台灣禮來股份有限公司
台北市復興北路365號11樓